

Confinement in the Three-dimensional Anisotropic t - J Model: A Mean-Field Study

Sanjoy K. Sarker

Department of Physics and Astronomy

The University of Alabama

Tuscaloosa, Alabama 35487

Abstract

We consider the anisotropic t - J model with the c-axis parameters t_c and J_c different from their in-plane counterparts, t and J . Within the slave-fermion mean-field approximation it is shown that the spiral state exhibits charge-confinement in the intermediate δ regime for a range of values of t_c/t . In the confined state the hopping amplitude $\langle c_{i\sigma}^\dagger c_{j\sigma} \rangle = 0$ along the c direction so that c-axis resistivity is infinite at $T = 0$.

The normal-state resistivity of the cuprate superconductors is metallic in the ab plane but is characteristic of an insulator along the c direction, leading to the possibility of the remarkable phenomenon of confinement [1, 2]. By continuity, such a behavior is not expected to occur in a Fermi liquid. In addition, confinement is at the heart of the proposed pair-tunneling mechanism which is in essence a deconfining process [3]. Theoretical treatments so far have been focused on a collection of weakly coupled Hubbard chains [2]. In this paper we study confinement in a collection of Hubbard planes, more specifically in the anisotropic t - J model described by the Hamiltonian

$$H = - \sum_{ij} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} + \frac{1}{2} \sum_{ij} J_{ij} [S_i \cdot S_j - n_i n_j]. \quad (1)$$

Here the in-plane hopping parameter $t_{ab} \equiv t$ and the exchange interaction $J_{ab} \equiv J$ are in general different t_c and J_c , the corresponding quantities along the c axis. Not all the parameters are free since the t - J model is thought to be derived from an underlying Hubbard model ($J_{ij} = 4t_{ij}^2/U$) so that $J_c/J = (t_c/t)^2$. There are therefore three independent parameters: t/J , $\xi \equiv t_c/t$ and the hole density δ .

Confinement is presumably intimately connected with spin-charge separation. While model (1) is not exactly solvable, in two dimensions a number of approximate ground states have been proposed that exhibit spin-charge separation. Here we study the spiral states in the Schwinger-boson slave-fermion representation: $c_{i\sigma} = h_i^\dagger b_{i\sigma}$, where $h^\dagger$ creates a fermionic hole and $b_{i\sigma}$ destroys a bosonic spin [4]. We will impose the constraint $h_i^\dagger h_i + \sum_\sigma b_{i\sigma}^\dagger b_{i\sigma} = 1$ on the average. A mean-field decomposition leads to following hamiltonians:

$$H_h = 2 \sum_{ij} t_{ij} B_{ij} h_i^\dagger h_j \quad (2)$$

$$H_b = \sum_{ij\sigma} t_{ij} D_{ij} b_{i\sigma}^\dagger b_{j\sigma} - \sum_{ij\sigma} J_{ij} A_{ij} \sigma b_{i\sigma} b_{j-\sigma}. \quad (3)$$

Here $D_{ij} = \langle h_i^\dagger h_j \rangle$ is the average hopping amplitude. This is associated with ferromagnetic backflow $B_{ij} = \langle b_{i\sigma}^\dagger b_{j\sigma} \rangle$. And $A_{ij} = \frac{1}{2} \langle (b_{i\uparrow} b_{j\downarrow} - b_{i\downarrow} b_{j\uparrow}) \rangle$ represents the antiferromagnetic correlations associated with the exchange term. We have shown that in two dimensions this competition gives rise to [4, 5] an incommensurate spiral metallic state that evolves

continuously from the Neel state at $\delta = 0$ to a ferromagnetic state at large $t\delta/J$. The spiral state is favored over double spiral and canted states and is stabilized against phase separation and domain walls by Coulomb repulsion [5, 6].

In the present problem the in-plane amplitudes (D_{ab} , B_{ab} and A_{ab}) are in general different from those along the c directions (D_c , B_c and A_c). We have solved the mean-field equations numerically for various values of t_{ab}/J_{ab} , $\xi = t_c/t_{ab}$ and δ . One self-consistent solution is found to be the usual spiral metallic state which evolves continuously out of the Neel state. For this state all the mean-field amplitudes including D_c and B_c are nonzero and hence there is no confinement. In addition, the state with $D_c = 0$ (and hence $B_c = 0$) is also a self-consistent solution. This is clearly a direct consequence of spin-charge separation. Charge can propagate in the ab plane in this case ($D_{ab} \neq 0$). Hence D_c can be thought of as the order-parameter for deconfining transitions.

While spin-charge separation can lead to confinement, it does not guarantee that such a state is favored energetically. The ground-state energy is given by

$$E_G = 8tD_{ab}B_{ab} - 4JR_{ab} + 4t_cD_cB_c - 2J_cR_c, \quad (4)$$

where $R_{ij} = A_{ij}^2 - B_{ij}^2/2 + (1-\delta)^2/8$. In general D and B have opposite signs and $R > 0$. For physically interesting values of $t - J$, we find that the unconfined phase has a lower energy both at small and large δ . But, as shown in Fig. 1, for intermediate δ there is a region in the δ - t_c/t plane where the *confined* state is favored. Interestingly, for fixed t/J and δ , the mean-field state is unconfined at small t_c/t , and with increasing t_c/t there is a first-order transition to a confined state. This is because as t_c decreases, $J_c \propto t_c^2$ decreases more rapidly. This favors ferromagnetic alignment and deconfinement. These results are summarized in Fig. 2.

For δ not too large the hole hopping amplitude D in the isotropic 3-D case is given by $D_3(\delta) \approx -\delta + \frac{(6\pi^2)^{2/3}}{10}\delta^{5/3}$. In two dimensions, $D_2(\delta) \approx -\delta + \pi\delta^2/2$. Hence, the hopping energy *per bond* can be lower in two dimensions. When t_c and δ are not too small, the system can gain maximum exchange energy in the c direction and maximum kinetic energy in the ab plane by having $B_c = D_c = 0$.

To summarize, we have shown that spiral state in the anisotropic t - J model exhibits charge confinement. For a more realistic treatment one needs to include fluctuations that destroy long-range magnetic order and reconstruct the Luttinger-Fermi surface, as shown previously for the 2-D model [7]. Such fluctuations are likely to bring the region of stability of the confined state to smaller values of δ and t_c/t . Nonetheless, there are some interesting consequences of confinement in our simple theory. (1) In the confined state the magnetic correlations along the c direction is peaked at $Q_c = \pi$, while the correlations in the ab plane remains incommensurate. (2) The c -axis resistivity is strictly infinite at $T = 0$, since $\langle c_{i\sigma}^\dagger c_{j\sigma} \rangle = -D_{ij} B_{ij} = 0$ along the c direction. However, at finite T or frequency there will be an incoherent contribution to the conductivity. Such a contribution will be activated if there is a spin gap.

References

- [1] P. W. Anderson, preprint.
- [2] D. G. Clarke, S. P. Strong and P. W. Anderson, *Phys. Rev. Lett.* **72**, 3218 (1994).
- [3] S. Chakravarty, A. Sudbo, P. W. Anderson and S. P. Strong, *Science* **261**, 337 (1993).
- [4] C. Jayaprakash, H.R. Krishnamurthy, and S.K. Sarker, *Phys. Rev. B***40**, 2610 (1989).
- [5] S. K. Sarker, *Phys. Rev. B* **46**, 8617 (1992).
- [6] F. M. Hu, S. K. Sarker and C. Jayaprakash, *Phys. Rev. B* **50**, 17901 (1994).
- [7] S. K. Sarker, *Phys. Rev. B***46**, 8617 (1992).

Figure Captions

Fig. 1. Phase diagram in the δ - t_c/t plane for two $t/J = 3$ (squares) and $t/J = 5$ (diamonds). In each case, the confined state ($D_c = 0$) has a lower energy in the V-shaped region. The transition is first order.

Fig. 2. The hole-hopping amplitude $D_c/(-\delta)$ and the ordering wavevector along the c-direction $Q_z \equiv \frac{Q_c}{2\pi}$ (diamonds) vs t_c/t . Note that Q_c is essentially zero (ferromagnetic) in the unconfined phase. Close to the transition it acquires a spiral character, and at the transition jumps to π corresponding to antiferromagnetic correlations in the confined phase.